arXiv:1406.3394v1 [math.NT] 13 Jun 2014

A note on the l-fold Bailey Lemma and Mixed Mock Modular forms

Alexander E Patkowski

Abstract

In this paper we present a method for constructing multiple-sum q-series for what is known as Mixed Mock Modular forms.

1 Introduction

A pair of sequences $(\alpha_{n_1,n_2,\cdots n_l},\beta_{n_1,n_2,\cdots n_l})$ is said to be a l-fold Bailey pair [4] with respect to a_j, $j=1,2,\cdots,l$ if

$$\beta_{n_1,n_2,\cdots n_l}=\sum_{r_1\geq 0}^{n_1}\sum_{r_2\geq 0}^{n_2}\cdots\sum_{r_l\geq 0}^{n_l}\frac{\alpha_{r_1,r_2,\cdots r_l}}{\prod_{i=1}^{l}(a_iq;q)_{n_i+r_i}(q;q)_{n_i-r_i}}. \tag{1.1}$$

The case $l=1$, is the standard well-known Bailey pair first introduced in [6], and $(U;q)_n=(1-U)(1-Uq)\cdots(1-Uq^{n-1})$ [9].

Several authors [8, 12] have discussed a modular form referred to as a *Mixed Mock Modular* form, which is of the form $\sum_{i\leq n}M_im_i$, where M_i is a mock modular form, and m_i is a modular form. A mixed mock modular form may be viewed as a generalization of a mock modular form.

This paper is concerned with applying results for the pair of sequences that satisfy (1.1) to obtain multi-sum q-series, which can be expressed as a product of a Mock modular form and a modular form. See Zwegers [15] work for important material on mock theta functions in relation to indefinite theta functions, and see Warnaar's paper [13] for a nice account of the Bailey lemma.

2 Observations on the l-fold Bailey Lemma

The purpose of this section is to outline some observations on the l-fold extension of Bailey's lemma which will prove fruitful in our applications to mixed mock modular forms. First, we

recall the commonly stated relation between the two sequences in a Bailey pair (α_n, β_n),

$$\sum_{n\geq 0}^{\infty}(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n\beta_n = \frac{(aq/\rho_1)_\infty(aq/\rho_2)_\infty}{(aq)_\infty(aq/\rho_1\rho_2)_\infty}\sum_{n\geq 0}^{\infty}\frac{(\rho_1)_n(\rho_2)_n(aq/\rho_1\rho_2)^n\alpha_n}{(aq/\rho_1)_n(aq/\rho_2)_n}. \tag{2.1}$$

Putting $\rho_1 = q^{-N_1}$, and $\rho_2 = q^{-N_2}$, gives

$$\sum_{j\geq 0}\frac{q^{j^2}a^j\alpha_j}{(aq)_{N_1+j}(aq)_{N_2+j}(q)_{N_1-j}(q)_{N_2-j}} = \frac{1}{(aq)_{N_1+N_2}}\sum_{j\geq 0}\frac{q^{j^2}a^j}{(q)_{N_1-j}(q)_{N_2-j}}\beta_j. \tag{2.2}$$

Setting (2.2) equal to β_{N_1,N_2} gives us a new 2-fold Bailey pair. Iterating this idea along the l-fold Bailey lemma gives us the following result.

Theorem 1. *If $(\alpha_{n_1,n_2,\cdots n_l}, \beta_{n_1,n_2,\cdots n_l})$ is a l-fold Bailey pair relative to $a_j = a$, $j = 1, 2, \cdots, l$, then $(\alpha_{n_1,n_2,\cdots n_{2l}}, \beta_{n_1,n_2,\cdots n_{2l}})$ is a $2l$-fold Bailey pair where*

$$\alpha_{n_1,n_2,\cdots n_{2l}} = \begin{cases} q^{r_1^2+r_2^2+\cdots+r_l^2}a^{r_1+r_2+\cdots r_l}\alpha_{r_1,r_2,\cdots r_l}, & \text{if } n_1 = n_2 = r_1, n_3 = n_4 = r_2, \cdots n_{2l-1} = n_{2l} = r_l, \\ 0, & \text{otherwise,} \end{cases} \tag{2.3}$$

and

$$\beta_{n_1,n_2,\cdots n_{2l}} = \frac{1}{\prod_{i\geq 1}^{l}(aq)_{n_{2i-1}+n_{2i}}}\sum_{i_1,i_2,\cdots i_l\geq 0}\frac{q^{i_1^2+i_2^2+\cdots+i_l^2}a^{i_1+i_2+\cdots i_l}\beta_{i_1,i_2,\cdots i_l}}{(q)_{n_1-i_1}(q)_{n_2-i_1}\cdots(q)_{n_{2l-1}-i_l}(q)_{n_{2l}-i_l}}. \tag{2.4}$$

The type of Bailey pairs we have constructed in Theorem 1 allow us relate a $2l$-fold q-series with a l-fold q-series. The case $l = 1$ of Theorem 1 gives a 2-fold Bailey pair of the Joshi-Vyas type [11].

3 Multi-sum q-series and mixed mock modular forms

Here we will consider applications of Theorem 1 to mixed mock modular forms. In particular, we shall prove the following result for two third-order mock theta functions and one tenth order mock theta function.

Theorem 2. *We have,*

$$\sum_{j,n_1,n_2\geq 0}\frac{q^{j^2+n_1^2+n_2^2+j+n_1+n_2}(-1)^j}{(q)_{n_1-j}(q)_{n_2-j}(q)_{n_1+n_2+1}(q^2;q^2)_j} = \frac{(-q)_\infty}{(q)_\infty}\sum_{n\geq 0}\frac{q^{2n^2+2n}}{(-q)_{2n+1}}, \tag{3.1}$$

$$\sum_{j,n_1,n_2\geq 0}\frac{(-q)_{n_1}(-q)_{n_2}q^{j^2+n_1(n_1+1)/2+n_2(n_2+1)/2+j}(-1)^j}{(q)_{n_1-j}(q)_{n_2-j}(q)_{n_1+n_2+1}(q^2;q^2)_j}=\frac{(-q)_\infty(-q;q^2)_\infty}{(q)_\infty(q;q^2)_\infty}\sum_{n\geq 0}\frac{q^{n^2+n}}{(-q;q^2)_{n+1}},\tag{3.2}$$

$$\sum_{j,n_1,n_2\geq 0}\frac{(q;q^2)_{n_1}(q;q^2)_{n_2}q^{2j^2+n_1^2+n_2^2}(-1)^{n_1+n_2+j-1}(q^2;q^4)_{j-1}}{(q^2;q^2)_{n_1-j}(q^2;q^2)_{n_2-j}(q^2;q^2)_{n_1+n_2}(q^2;q^2)_{2j-1}}=\frac{(q;q^2)_\infty^2(q^4;q^4)_\infty}{(q^2;q^2)_\infty^2(-q^4;q^4)_\infty}\psi(q^4),\tag{3.3}$$

where $\psi(q)=\sum_{n\geq 1}q^{n(n+1)/2}/(q;q^2)_n$.

Proof. First recall [4] that

$$\sum_{n_1\geq 0}^{\infty}\sum_{n_2\geq 0}^{\infty}(x)_{n_1}(y)_{n_1}(z)_{n_2}(w)_{n_2}(a_1q/xy)^{n_1}(a_2q/zw)^{n_2}\beta_{n_1,n_2}$$

$$=\frac{(a_1q/x)_\infty(a_1q/y)_\infty(a_2q/z)_\infty(a_2q/w)_\infty}{(a_1q)_\infty(a_1q/xy)_\infty(a_2q)_\infty(a_2q/zw)_\infty}\sum_{n_1\geq 0}^{\infty}\sum_{n_2\geq 0}^{\infty}\frac{(x)_{n_1}(y)_{n_1}(z)_{n_2}(w)_{n_2}(a_1q/xy)^{n_1}(a_2q/zw)^{n_2}\alpha_{n_1,n_2}}{(a_1q/x)_{n_1}(a_1q/y)_{n_1}(a_2q/z)_{n_2}(a_2q/w)_{n_2}}.\tag{3.4}$$

The case (3.1): Insert the Bailey pair (α_n,β_n) relative to $a=q$,

$$\alpha_n=\frac{q^{n^2}(1-q^{2n+1})}{(1-q)}\sum_{|j|\leq n}(-1)^jq^{-j^2},\tag{3.5}$$

$$\beta_n=\frac{(-1)^n}{(q^2;q^2)_n},\tag{3.6}$$

into (2.2) and insert the resulting 2-fold pair

$$\alpha_{N_1,N_2}=\begin{cases}\frac{q^{2n^2+n}(1-q^{2n+1})}{(1-q)}\sum_{|j|\leq n}(-1)^jq^{-j^2}, & \text{if } N_1=N_2=n\\ 0, & \text{otherwise,}\end{cases}\tag{3.7}$$

and

$$\beta_{N_1,N_2,\cdots N_{2l}}=\frac{1}{(q)_{N_1+N_2+1}}\sum_{j\geq 0}\frac{q^{j^2+j}(-1)^j}{(q)_{N_1-j}(q)_{N_2-j}(q^2;q^2)_j}.\tag{3.8}$$

into (3.4) with $x,y,z,w\to\infty$, $a_1=a_2=q$ to get

$$\sum_{j,n_1,n_2\geq 0}\frac{q^{j^2+n_1^2+n_2^2+j+n_1+n_2}(-1)^j}{(q)_{n_1-j}(q)_{n_2-j}(q)_{n_1+n_2+1}(q^2;q^2)_j}=\frac{1}{(q)_\infty^2}\sum_{n\geq 0}q^{4n^2+3n}(1-q^{2n+1})\sum_{|j|\leq n}(-1)^jq^{-j^2}.\tag{3.9}$$

Note that the indefinite quadratic series on the right side of (3.9) is found to be related to a mock theta function noted by Andrews by [5, eq.(1.15)]. This completes the proof of (3.1).

The case (3.2): First we note a simple consequence of the second pair in table one in [2, pg.75] inserted in a limiting case of Bailey's lemma ($\rho_1=-q$, $\rho_2\to\infty$, in (2.1)),

$$\sum_{n\geq 0}\frac{q^{n(n+1)}}{(-q;q^2)_{n+1}}=\frac{(-q^2;q^2)_\infty}{(q^2;q^2)_\infty}\sum_{n\geq 0}q^{3n^2+2n}(1-q^{2n+1})\sum_{|j|\leq n}(-1)^jq^{-j^2}.\tag{3.10}$$

(The q-series on the left side is a third-order mock theta function [14].) Inserting (3.7)–(3.8) into (3.4) with $x = z = -q, y, w \to \infty$, $a_1 = a_2 = q$, gives the result.

The case (3.3): Inserting the pair due to Bringmann and Kane [7, Theorem 2.3, (1)] into (2.2) and then inserting the resulting pair into (3.4) with $x = z = -q^{1/2}, y, w \to \infty$, $a_1 = a_2 = 1$, gives

$$\sum_{j,n_1,n_2\geq 0} \frac{(-q^{1/2})_{n_1}(-q^{1/2})_{n_2}q^{j^2+n_1^2/2+n_2^2/2}(-1)^j(q;q^2)_{j-1}}{(q)_{n_1-j}(q)_{n_2-j}(q)_{n_1+n_2}(q)_{2j-1}}$$

$$= \frac{(-q^{1/2})_\infty^2}{(q)_\infty^2}\left(\sum_{n\geq 1} q^{10n^2-2n}(1-q^{4n})\sum_{j=-n}^{n-1} q^{-2j^2-2j} - \sum_{n\geq 0} q^{10n^2+8n+2}(1-q^{4n+2})\sum_{|j|\leq n} q^{-2j^2}\right). \tag{3.11}$$

On the other hand, the work of Choi [9] gives us

$$\sum_{n\geq 1}\frac{q^{n(n+1)/2}}{(q;q^2)_n} = \frac{(-q)_\infty}{(q)_\infty}\left(\sum_{n\geq 0} q^{5n^2+4n+1}(1-q^{2n+1})\sum_{|j|\leq n} q^{-j^2} - \sum_{n\geq 1} q^{5n^2-n}(1-q^{2n})\sum_{j=-n}^{n-1} q^{-j^2-j}\right). \tag{3.12}$$

This follows from the pair [1, pg.131, eq.(7.18)–(7.19)] inserted into the $\rho_1 = -1$, $\rho_2 \to \infty$ case of (2.1). Hence comparing the indefinite theta functions in (3.11) and (3.12) and a little manipulation gives (3.3).

□

4 More Comments

We mention a few more notes on our work. First, it should be noted the observation made in the second section may be applied in other directions. For example, the identity (2.2) may be used to obtain many interesting q-polynomial identities. The choice $\alpha_0 = 1$, $\alpha_n = (-1)^n q^{n(3n-1)/2}(1+q^n)$, $n > 0$, and $\beta_n = 1/(q)_n$, gives us a key identity used in [3, eq.(5.33)], which may in turn be used to obtain [3, eq.(5.30)] after appealing to the $l = 2$ case of Theorem 1. We leave the details to the reader. A direct corollary of this remark leads us to the Rogers-Ramanujan type identity (by using [4, Corollary 1, $s = 2$])

$$\sum_{j,n_1,n_2\geq 0}\frac{q^{j^2+n_1^2+n_2^2}}{(q)_{n_1+n_2}(q)_{n_1-j}(q)_{n_2-j}(q)_j} = \frac{(q,q^8,q^9;q^9)_\infty}{(q)_\infty^2}. \tag{4.1}$$

Second, to see how one may apply Theorem 1 in its full generality, we may consider the pair $\alpha_0 = 1$, 0 otherwise, and $\beta_n = 1/(q)_n^2$. Using this pair with (2.2) gives us

$$\frac{1}{(q)_{n_1}^2(q)_{n_2}^2} = \frac{1}{(q)_{n_1+n_2}}\sum_{j\geq 0}\frac{q^{j^2}}{(q)_{n_1-j}(q)_{n_2-j}(q)_j^2}, \tag{4.2}$$

which is equivalent to the 2-fold Bailey pair

$$\alpha_{n_1,n_2} = \begin{cases} 1, & \text{if } n_1 = n_2 = 0, \\ 0, & \text{otherwise,} \end{cases} \tag{4.3}$$

and

$$\beta_{n_1,n_2} = \frac{1}{(q)_{n_1+n_2}} \sum_{j\geq 0} \frac{q^{j^2}}{(q)_{n_1-j}(q)_{n_2-j}(q)_j^2}. \tag{4.4}$$

Applying this pair to the $l = 2$ case of Theorem 1 gives us

$$\alpha_{n_1,n_2,n_3,n_4} = \begin{cases} 1, & \text{if } n_1 = n_2 = n_3 = n_4 = 0, \\ 0, & \text{otherwise,} \end{cases} \tag{4.5}$$

and

$$\beta_{n_1,n_2,n_3,n_4} = \frac{1}{(q)_{n_1+n_2}(q)_{n_3+n_4}} \sum_{i_1,i_2,j\geq 0} \frac{q^{i_1^2+i_2^2+j^2}}{(q)_{n_1-i_1}(q)_{n_2-i_1}(q)_{n_3-i_2}(q)_{n_4-i_2}(q)_{i_1+i_2}(q)_{i_1-j}(q)_{i_2-j}(q)_j^2}. \tag{4.6}$$

Therefore we may use the l-fold Bailey lemma to give us

$$\sum_{n_1,n_2,j\geq 0} \frac{q^{j^2+n_1^2+n_2^2}}{(q)_{n_1+n_2}(q)_{n_1-j}(q)_{n_2-j}(q)_j^2} = \frac{1}{(q)_\infty^2}, \tag{4.7}$$

and

$$\sum_{n_1,n_2,n_3,n_4,i_1,i_2,j\geq 0} \frac{q^{n_1^2+n_2^2+n_3^2+n_4^2+i_1^2+i_2^2+j^2}}{(q)_{n_1+n_2}(q)_{n_3+n_4}(q)_{n_1-i_1}(q)_{n_2-i_1}(q)_{n_3-i_2}(q)_{n_4-i_2}(q)_{i_1+i_2}(q)_{i_1-j}(q)_{i_2-j}(q)_j^2} = \frac{1}{(q)_\infty^4}. \tag{4.8}$$

Continuing this idea leads us to an expression for $1/(q)_\infty^{2M}$, $M \in \mathbb{N}$.

References

[1] G.E. Andrews, *The fifth and seventh order mock theta functions,* Trans. Amer. Math. Soc. 293 (1986), no. 1, 113–134.

[2] G.E. Andrews and D. Hickerson, *Ramanujan's "lost" notebook. VII. The sixth order mock theta functions,* Adv. Math. 89 (1991), no. 1, 60–105.

[3] G.E. Andrews, A. Schilling, and S.O. Warnaar, *An A2 Bailey lemma and Rogers–Ramanujan-type identities,* Journal of the American Mathematical Society 12 (1999), 677–702.

[4] George E. Andrews, *Umbral Calculus, Bailey Chains, and Pentagonal Number Theorems,* J. Comb. Theory, Ser. A 91 (1-2): 464–475 (2000)

[5] George E. Andrews, *q-Orthogonal polynomials, Rogers-Ramanujan identities, and mock theta functions,* Proceedings of the Steklov Institute Dedicated to the 75th Birthday of A. A. Karatsuba, 276:21-32 (2012)

[6] W. N. Bailey, *Identities of the Rogers–Ramanujan type*, Proc. London Math. Soc. (2), 50 (1949), 1–10.

[7] K. Bringmann and B. Kane, *Multiplicative q-hypergeometric series arising from real quadratic fields,* Trans. Amer. Math. Soc. 363 (2011), no. 4, 2191–2209.

[8] K. Bringmann and B. Kane, *Second order cusp forms and mixed mock modular forms,* Ramanujan Journal, special volume in honour of Ismail and Stanton 31, 147-161 (2013).

[9] Y.-S. Choi, *Tenth order mock theta functions in Ramanujan's Lost Notebook,* Invent. Math. 136 (1999), 497–569.

[10] G. Gasper, M. Rahman, *Basic hypergeometric series*, Cambridge Univ. Press, Cambridge, 1990.

[11] C.M. Joshi and Y. Vyas, *Bailey Type Transforms and Applications* http://arxiv.org/abs/math/0701192

[12] J. Lovejoy and R. Osburn, *Mixed mock modular q-series,* J. Indian Math. Soc., Special Volume to commemorate the 125th Birth Anniversary of Srinivasa Ramanujan and the National Mathematics Year - 2012 (2013), 45-61.

[13] S. O. Warnaar, *50 years of Bailey's lemma,* Algebraic combinatorics and applications (Goweinstein, 1999), 333–347, Springer, Berlin, 2001.

[14] G. N. Watson, *The final problem: an account of the mock theta functions,* J. London Math. Soc., 11(1936), 55–80.

[15] S. Zwegers, *Mock Theta Functions,* PhD Thesis, Universiteit Utrecht (2002).

1390 Bumps River Rd.

Centerville, MA 02632

USA

E-mail: alexpatk@hotmail.com